Problem Set 7

Discrete Mathematics

Due on the 27^{th} of March, 2024

(10 pts)   1. Let X be a set. Show that $(\forall Y \in \mathbb{P}(X))\,(|Y| \leqslant |X|)$.

(15 pts)   2. Show that $\forall X \forall Y \Big(|X| \leqslant |Y| \Rightarrow \exists Z \big(Z \subseteq Y \wedge |X| = |Z| \big) \Big)$.

(15 pts)   3. Let X, Y, Z be sets and consider $f : X \to Y$ and $g : Y \to Z$. We define the *composition* of f with g to be the function $g \circ f : X \to Z$ given by $(g \circ f)(x) := g(f(x))$ for all $x \in X$.

Since the codomain of f and the domain of g are the same, they are *compatible*, and their composition is sensibly defined.

   (a) Show that, if f and g are both injections, then $g \circ f$ is injective.

These are called *monomorphisms*.

   (b) Show that, if f and g are both surjections, then $g \circ f$ is surjective.

These are called *epimorphisms*.

   (c) Show that, if f and g are both bijections, then $g \circ f$ is bijective.

These are called *isomorphisms*.

(30 pts)   4. For this problem, let X and Y be nonempty sets and let $f : X \to Y$.

   (a) If f is injective, show there exists $g : Y \to X$ where $g \circ f = \text{id}_X$.

   (b) If f is surjective, show there exists $g : Y \to X$ where $f \circ g = \text{id}_Y$.

   (c) If f is a bijection, then show that there exists a ~~unique~~ function $g : Y \to X$ such that $g \circ f = \text{id}_X$ and $f \circ g = \text{id}_Y$.

(30 pts)   5. *Euler's totient function* is the function $\varphi_e : \mathbb{N} \to \mathbb{N}$ that counts how many positive integers are *coprime* with each $n \in \mathbb{N}$, defined below.

$$\varphi_e(n) := \Big| \big\{ z \in \mathbb{N} \mid 1 \leqslant z \leqslant n \wedge \gcd(z, n) = 1 \big\} \Big|$$

   (a) If $p, k, m \in \mathbb{N}_+$ are *positive* naturals with p prime and $m \leqslant p^k$, then prove that $\gcd(p^k, m) \neq 1 \Leftrightarrow p \mid m$.

   (b) If p is prime, then prove that $\varphi_e(p) = p - 1$.

   (c) If p is prime and $k \in \mathbb{N}_+$, then prove that $\varphi_e(p^k) = p^k - p^{k-1}$.

Hint: count the multiples of p.